84-4145

FISHERY SURVEY AND RELATED LIMNOLOGICAL CONDITIONS OF WILLIAMS LAKE, HUBBARD COUNTY, MINNESOTA

U.S. GEOLOGICAL SURVEY

Water-Resources Investigations Report 84-4145

NEWARK PUBLIC LIBRARY

MAR 5 1986

U. S. GOVT. PUBS.
DEPOSITORY 376

I 19.42/4:

READEX MICROPRINT best copy available

FISHERY SURVEY AND RELATED LIMNOLOGICAL CONDITIONS OF WILLIAMS LAKE, HUBBARD COUNTY, MINNESOTA

By William W. Taylor, James W. LaBaugh, Mark H. Freeberg and David C. Dowling

U.S. GEOLOGICAL SURVEY

Water-Resources Investigations Report 84-4145

Lakewood, Colorado
1985

UNITED STATES DEPARTMENT OF THE INTERIOR

DONALD PAUL HODEL, Secretary

GEOLOGICAL SURVEY

Dallas L. Peck, Director

For additional information
write to:

James W. LaBaugh
U.S. Geological Survey,
Box 25046, MS 413
Denver Federal Center
Lakewood, CO 80225

Copies of this report can
be purchased from:

Open-File Services Section
Western Distribution Branch
U.S. Geological Survey, MS 306
Box 25425, Federal Center
Denver, CO 80225
Telephone: (303) 236-7476

CONTENTS

	Page
Abstract	1
Introduction	1
Methods	2
Limnological conditions	2
Fishery survey	5
Species composition	5
Population characteristics	5
Food habits	13
Summary	13
References cited	25

ILLUSTRATIONS

		Page
Figure 1.	Graphs showing vertical distribution of selected limnological characteristics, August 27, 1982	3
2.	Map showing the extent of the distribution of emergent, floating-leaved, and submerged aquatic macrophytes, July-August 1982	6
3.	Map showing relative distribution of benthic invertebrates in the littoral zone, 1980-82	8
4-10.	Graphs showing population characteristics of fish collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight:	
4.	Bluegill	10
5.	Pumpkinseed sunfish	12
6.	Largemouth bass	15
7.	Northern pike	17
8.	Yellow perch	19
9.	Rock bass	21
10.	Black crappie	23

TABLES

		Page
Table 1.	Composition of the zooplankton community, August 27, 1982-------	4
2.	Taxa of benthic invertebrates in the littoral zone, 1980-82-----	7
3.	Species of fish captured, August 31-September 2, 1982----------	9
4-10.	Mean back-calculated lengths for fish collected in August 1982:	
4.	Bluegill----------	9
5.	Pumpkinseed----------	11
6.	Largemouth bass----------	14
7.	Northern pike----------	16
8.	Yellow perch----------	18
9.	Rock bass----------	20
10.	Black crappie----------	22
11.	Stomach analysis of fish collected, August 31-September 2, 1982----------	24

CONVERSION FACTORS

For the convenience of those readers who prefer to use inch-pound units rather than the International System of Units (SI), the conversion factors for units used in this report are listed below:

Multiply SI unit	*By*	*To obtain inch pound unit*
centimeter (cm)	0.393	inch
meter (m)	3.281	feet
siemens (S)	1.000	mhos

FISHERY SURVEY AND RELATED LIMNOLOGICAL CONDITIONS OF WILLIAMS LAKE, HUBBARD COUNTY, MINNESOTA

By William W. Taylor[1], James W. LaBaugh[2], Mark H. Freeberg[1], and David C. Dowling[1]

ABSTRACT

Bluegill (*Lepomis macrochirus*), pumpkinseed sunfish (*Lepomis gibbosus*), largemouth bass (*Micropterus salmoides*), yellow perch (*Perca flavescens*), rock bass (*Amploplites rupestris*), black crappie (*Pomoxis nigromaculatus*), and northern pike (*Esox lucius*) were found in Williams Lake, Hubbard County, Minnesota, during a fishery survey of the lake in late August 1982. The most abundant fish were the bluegills. These fish live in the large littoral zone of the lake; this zone underlied 55 percent of the surface area of the lake. The most ubiquitous benthic invertebrate in the littoral zone (*amphipods*) and the most abundant benthic invertebrate (*chironomid larvae*) were major food items for the bluegill. Other organisms found in the stomach contents of fish collected in this survey were zooplankton, gastropods, *Diptera* larvae, odonates, terrestrial insects, and other fish. Daphnia were the only zooplankters of a diverse plankton community that were found in stomach contents. The abundance of fish other than bluegill was typical for a system in which northern pike is the major predator.

INTRODUCTION

Williams Lake, a small lake in north-central Minnesota, is one of several natural lakes selected by the U.S. Geological Survey and its cooperators for intensive field research on the interactions of the hydrologic cycle and nutrient balances of lakes (Siegel and Winter, 1980; LaBaugh and others, 1981). One of the objectives of the ongoing project at this lake is to interrelate the physical and chemical characteristics of the water to biological production. In a preliminary effort to evaluate the fish community in the lake, an exploratory survey of the fish population was conducted in late August 1982. The objectives of this survey were to ascertain species composition, their vital statistics, and their food habits for future reference of the potential impact of fishes in the nutrient cycling and energy flow of this lake. The purpose of this report is to place the findings of the fishery survey in the context of limnological conditions in Williams Lake.

[1]Department of Fisheries and Wildlife, Michigan State University, East Lansing, Michigan.

[2]U.S. Geological Survey, Denver, Colorado.

METHODS

Samples for chemical and plankton community analyses were collected by using methods described by LaBaugh and others (1981). Benthic samples were collected along six transects in the near-shore area of the littoral zone of the lake. Samples were collected by pushing a 20-cm diameter plastic coring tube at least 5 cm into the bottom material. Contents of the tube were held in place with a piston from a Livingstone piston corer. Next, the tube was removed from the bottom sediments. Then, the contents were emptied into plastic containers and formalin was added to preserve the sample for subsequent analysis. The distance from shore at which the zone of emergent, floating-leaved, and submerged aquatic macrophytes ended was measured along the same transects along which benthic samples were collected. In between transects, the limit of occurrence for submerged macrophytes was inferred from the depth contour map of the lake. Benthic samples were collected in July 1980, October 1981, and July 1982. The survey of the aquatic macrophytes in Williams Lake occurred in July-August 1982.

Collection of fish began on August 31, 1982, and continued through September 2, 1982, using fyke nets and gill nets in accordance with provisions in a permit issued by the Minnesota Department of Natural Resources. Electrofishing techniques were tried but were ineffectual due to the abundance of aquatic vegetation in the littoral zone. Four 2.54-cm stretch-mesh fyke nets with a 15.2-m lead were used during the 48-hour survey. Net placement was planned after a preliminary examination of the physical characteristics of the littoral zone of the lake. The basic procedure was to choose a site randomly for net placement in each observably different habitat type--vegetated, rocky, or sandy. In addition, an attempt was made to collect fish in shallow (<1.5-m depth) and deep water. Samples from the fyke nets were collected after 12 hours at each sampling site; the cumulative collection time was 96 hours.

Two experimental gill nets were used in the littoral and limnetic regions of Williams Lake. The nets were 150 ft in length and were composed of six 25-ft panels. The mesh sizes of the stretch-mesh panels were: 2.54 cm, 3.81 cm, 5.08 cm, 6.36 cm, 7.62 cm, and 10.16 cm. Captured fish were identified to species and measured for length and weight. Additionally, several body scales were removed from selected fish for age-growth analysis. The fish were then released, live, to Williams Lake.

A preliminary analysis of stomach contents was performed on one to four fish of each captured species, except northern pike. The number of fish sacrificed was limited by the provisions of the collecting permit issued by the Minnesota Department of Natural Resources.

LIMNOLOGICAL CONDITIONS

A summary of the vertical distribution of temperature, specific conductance, pH, and dissolved oxygen, measured on August 27, 1982, is shown in figure 1. The lake was thermally stratified. An inverse clinograde vertical distribution was measured for specific conductance. A clinograde vertical distribution was measured for dissolved oxygen and pH. These conditions in


DEPTH BELOW LAKE SURFACE, IN METERS
0
5
9
0 5 10 15 20 25 30 35 40
TEMPERATURE, IN DEGREES CELSIUS
0 50 100 150 200 250 300 350 400
SPECIFIC CONDUCTANCE,
IN MICROSIEMENS PER CENTIMETER AT 25°CELSIUS
4 5 6 8 9 10 11 12 13 14
PH
PERCENT SATURATION OF DISSOLVED OXYGEN
0 50 100 150 200
Concentration
percent saturation
0 5 10 15 20
DISSOLVED-OXYGEN CONCENTRATION, IN MILLIGRAMS PER LITER

Figure 1.--Vertical distribution of selected limnological characteristics, August 27, 1982.

the water column were similar to those reported for August 1979 by LaBaugh and others (1981) and for the same time in 1980 and 1981 (based on unpublished data). The Secchi-disk transparency of 4.8 m measured on August 27, 1982, also was typical for late August.

LaBaugh and others (1981) indicated blue-green algae and copepods were the dominant groups of the planktonic community of Williams Lake, when it was thermally stratified in the summer of 1979. At the time of the fish population survey reported here, blue-green algae again were the numerically dominant group among the phytoplankton. However, the most abundant zooplankton group was the rotifers (table 1). *Daphnia galeata mendotae* was the only species of Daphnia present in the zooplankton community. Daphnia were the only zooplankters found in the stomach contents of fish collected during this investigation.

Table 1.--*Composition of the zooplankton community, August 27, 1982*

Taxa	Number of organisms per liter	
	Replicate 1	Replicate 2
Copepoda	50.1	50.6
Orthocyclops modestus	----	1.8
Cyclops biscuspidatus thomasi	1.4	2.8
Cyclopoid copepodids	11.6	9.3
Diaptomus oregonensis	----	1.0
Calanoid copepodids	7.0	3.1
Copepod naupli	30.1	32.6
Cladocera	60.3	61.5
Diaphanosoma leuctenbergianum	7.4	8.0
Daphnia galeata mendotae	6.5	8.5
Ceriodaphnia lacustris	.5	----
Bosmina longirostris	.5	1.0
Chydorus spaericus	45.4	44.0
Rotifera	55.6	95.1
Conochilus unicornis	19.9	45.8
Polyarthra dolichoptera	.5	----
Polyarthra vulgaris	----	1.0
Gastropus hyptopus	.9	1.0
Asplanchna priodonta group	.5	1.3
Kellicottia longispina	4.6	4.4
Keratella cochlearis	26.4	36.2
Monostyla sp.	1.4	4.4
Trichocerca sp	1.4	1.0

In July and August of 1982, a preliminary survey was made of the distribution of aquatic macrophytes in Williams Lake; results of that survey are shown in figure 2. The area of the lake surface containing or underlain by aquatic macrophytes was 55 percent of the total surface area. Submersed macrophytes were found throughout the littoral zone. Floating-leaved macrophytes were absent from most of the northeast side of the lake. Sampling of the fish population with fyke nets was designed to avoid any possible bias introduced by the spatial variability of the aquatic-macrophyte community.

The benthic-invertebrate community of the littoral zone was diverse (table 2). The amphipod *Hyatella azteca* was a ubiquitous member of the benthic community, being found at every sampling transect during the October 1981 collection. Amphipods were the most numerous prey found among the stomach contents of bluegills. Chironomids usually were the most abundant organisms at each sampling transect (fig. 3). Chironomids were also prey of bluegills.

FISHERY SURVEY

Species Composition

Seven species were captured during this fishery investigation. The predominant species in Williams Lake is the bluegill (*Lepomis macrochirus*), which lives in the vegetation along the shore line (table 3). The pumpkinseed sunfish (*Lepomis gibbosus*) and largemouth bass (*Micropterus salmoides*) would be considered common, with northern pike (*Esox lucius*) present in average abundance for a top carnivore. Yellow perch (*Perca flavescens*), rock bass (*Amploplites rupestris*), and black crappie (*Pomoxis nigromaculatus*) were captured infrequently.

Population Characteristics

The majority of bluegills collected were between 12 and 16 cm in total length (fig. 4*A*). These fish primarily were age 4 and 5 years (fig. 4*B*). These fish were smaller than expected for a normal population (Carlander, 1977) indicating that an overabundance of bluegill occurs, which inhibits their growth (table 4).

The length of fish in any year of life prior to collection of a scale for growth analysis is shown for bluegill from Williams Lake in table 4. The method of comparing scale size (radius) to fish size (length) at a known age, and then determining what length the fish must have been at an earlier age, is known as back calculation. For example, bluegill sampled in this study whose scale annuli indicated they were 3 years old or older, but less than 4 years old, had a mean length of 10.7 cm. Fish in this age group were not all exactly 3 years old, so the length of fish at age 3 will be some value less than 10.7; some growth will have occurred in those fish older than 3 years in age. By comparing scale growth to fish length, the average length of bluegill fish at 3 years of age was calculated to be 10.4 cm. When these 3-year old fish were 1 year old, their average length was calculated to be 7.3 cm. These data are used to interpret the growth characteristics of the fish population in the lake.




95°40'
EXPLANATION
MACROPHYTE DISTRIBUTION
From transects A, B, C, D, E, H, X, and G
AREA OF MACROPHYTE GROWTH --
Line is dashed where boundary is inferred
AREA OF FLOATING-LEAF
MACROPHYTE GROWTH
46°57'30"
A
TRANSECT AND LABEL--Emergent macrophytes found at A, B, C, D, G at 0 to 5 meters from shore. Emergent macrophytes not found at E, H, and X
10
LINE OF EQUAL DEPTH OF WATER --
Interval is 2 feet. Datum is lake surface
12
0
300 METERS
0
900
1000 FEET

Figure 2.--Extent of the distribution of emergent, floating-leaved, and submerged aquatic macrophytes, July-August 1982.

Table 2.--*Taxa of benthic invertebrates in the littoral zone, 1980-82.*

Diptera

Chironomidae

Cladotanytarsus *Trissocladium* *Cryptochiromus* *Zaurilimyia*
Psectrocladius *Trichocladius* *Tanytarsus* *Labrundinea*
Stichtochiromomus *Diamesa* *Procladius* *Dicrotendipes*
Larsia *Paratanytarsus* *Lauterborniella* *Pseudochironomus*
Polypedilum *Phaenospectra* *Microtendipes* *Psectrotanypus*

Ceraptogonidae

Bezzia, Probezzia *Stilobezzia*

Tabanidae

Chrysops

Palpomyia *Culicoides*

Trichoptera

Molanna
Phryganea
Oxythria
Oecetis
Ceraclea
Mystacides

Ephemeroptera

Choroterpes
Paracloedes
Caenis
Hexagenia
Ephemerella

Odonata

Gomphus
Enallagma
Libellula

Amphipoda

Hyatella azteca

Gastropoda

Amnicola
Gyraulus
Physa

Coleoptera

Donacia
Halipus

Pelecypoda

Limnodrilus
Pisidium

Oligocheata

Chaetogaster limnaei
Specaris josinae
Pristina
Peloscolex
Tubificidae
Nais communis
Dero digitata
Amphicheata americanus
Lumbriculidae

Spaherium

Stylarsia fossularis
Stylaria lacustris
Stylodrilus heringianus
Chaetogaster cristallinus
Lumbriculus variagatus
Homochaeta
Dero nivca
Aulodrilus
Uncinais uncinata

Nematomorpha

Gordius

Polychaeta

Manyonkia speciosa

Nematoda

Bactrachobdella
Eudorylaimus
Tylenchus
Aphanolaimus

Hirudinea

Arrenurus

Hydracarina

Anonchus



Figure 3.--Relative distribution of benthic invertebrates in the littoral zone, 1980-82.

Table 3.--*Species of fish captured, August 31-September 2, 1982*

[Fish listed in order of relative abundance from most to least]

Species name	Common name
Lepomis macrochirus	bluegill
Lepomis gibbosus	pumpkinseed sunfish
Micropterus salmoides	largemouth bass
Esox lucius	northern pike
Perca flavescens	yellow perch
Ambloplites rupestris	rock bass
Pomoxis nigromaculatus	black crappie

Table 4.--*Mean back-calculated lengths for bluegill collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
1	6.9	3	6.7						
2	8.1	8	6.9	8.4					
3	10.7	5	7.3	8.8	10.4				
4	13.5	21	7.9	9.6	11.1	12.5			
5	14.2	31	7.4	9.0	10.4	11.3	12.6		
6	15.9	8	7.5	9.1	10.8	11.7	12.7	13.7	
7	16.9	2	7.3	8.4	9.7	10.8	11.7	12.4	13.3
Mean length of fish at each age			7.3	8.9	10.4	11.6	12.3	13.0	13.3
Annual mean growth increment			1.6	1.5	1.2	.7	.7	.2	

*Calculated average length of fish for each age class at a specific age in their life history.


A
Number of fish = 200
PERCENT
LENGTH AT MIDDLE OF INTERVAL, IN CENTIMETERS
B
Number of fish = 78
PERCENT
AGE, IN YEARS
C
Number of fish = 146
Regression equation Log weight = -1.77 + 2.97 × Log length
R² = .94
DATA POINTS--Each symbol represents the corresponding number of data points
LOG WEIGHT, IN GRAMS
LOG LENGTH, IN CENTIMETERS

Figure 4.--Population characteristics of bluegill collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Growth characteristics of fish populations also can be examined by use of the relation between weight and length. The weight-length relationship can be expressed as:

$$W = aL^b, \quad (1)$$

where

W is weight of the fish;
L is length of the fish;
a is an intercept value; and
b is an exponent, indicating isometric or allometric growth and its rate (Ricker, 1975).

Generally, a "b" value of 3 indicates a fish is growing isometrically; body proportions are unchanged with growth (Bagenal, 1978). A value greater or less than 3 indicates allometric growth; that is, the fish becomes heavier or lighter for its length as it grows longer and body proportions change with growth. Bluegills in Williams Lake are growing nearly isometrically as indicated by a "b" value of 2.97 (fig. 4*C*), which is slower than a normally growing population (Carlander, 1977).

Forty-three percent of the pumpkinseed sunfish collected were 16 cm in total length (fig. 5*A*). These fish primarily were 3 years old (fig. 5*B*). Growth was slightly greater than average compared with those reported by other investigators (Carlander, 1977)(table 5, fig. 5*C*).

Table 5.--*Mean back-calculated lengths for pumpkinseed sunfish collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
2	9.4	5	8.2	9.5					
3	11.1	6	7.9	8.7	10.0				
5	15.3	3	8.5	9.7	11.1	12.1	13.3		
6	16.6	1	10.1	11.6	12.7	14.3	15.2	16.0	
Mean length of fish at each age----------			8.7	9.9	11.2	13.2	14.2	16.0	
Annual mean growth increment----------			1.2	1.3	2.0	1.0	1.8		

*Calculated average length of fish for each age class at a specific age in their life history.



Figure 5.--Population characteristics of pumpkinseed sunfish collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

The largemouth bass population had two dominant size classes, one at 14 cm and one at 26 cm (fig. 6*A*). Overall, 4-year-old bass predominated, comprising 32 percent of the catch (fig. 6*B*). The length-weight and growth relationship appeared to be average for this species (Carlander, 1977) (table 6, fig. 6*C*).

Northern pike collected in the survey ranged between 38 and 78 cm in length (fig. 7*A*). Because only 10 fish were captured, our analysis of their vital statistics is incomplete. However, data for the age, length, weight, and growth relationships are given in table 7 and figure 7*B,C*. Caution needs to be used in the interpretation and extrapolation of these data in relation to the total population because of the small sample size. The same is true for yellow perch (table 8, fig. 8*A-C*), rock bass (table 9, fig. 9*A-C*), and black crappie (table 10, fig. 10*A-C*) populations.

Food Habits

Stomach analyses of selected species are depicted in table 11. Bluegills fed primarily on amphipods during this study; they, in turn, were eaten by largemouth bass, rock bass, and black crappie. Largemouth bass had the greatest diversity in diet, with some fish eating from the water mass (bluegills, *Daphnia*), and others feeding on the bottom organisms (Amphipods, Chironomid larvae, Diptera larvae, Odonata). Northern pike were not examined for food studies because of their abundance and size.

SUMMARY

It is clear from this study that the bluegill is a key component in the nutrient and biological energy budget of Williams Lake. Bluegills are abundant (overabundant) living in the very productive littoral zone and feeding on the benthic dwelling invertebrates. These fish, in turn, provide food for the piscivorous fishes, which live in the limnetic and shoreline zone of this lake. Further research, centering on the role of the bluegill in the energy flow and nutrient cycling in Williams Lake, is needed to assess the effect of fish in the functional development of this ecosystem.

Table 6.--*Mean back-calculated lengths for largemouth bass collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
1	6.5	2	5.3						
2	13.6	5	7.0	11.6					
3	21.3	6	7.6	13.1	18.7				
4	25.4	9	6.5	11.6	16.0	21.3			
5	28.6	4	6.8	10.3	15.1	19.6	24.1		
6	31.5	1	6.7	10.5	13.8	19.6	22.7	27.7	
8	43.2	11	5.4	11.9	17.6	23.9	30.2	34.8	39.2	42.5
Mean length of fish at each age----------			6.5	11.5	16.2	21.1	25.7	31.3	39.2	42.5
Annual mean growth increment----------			5.0	4.7	4.9	4.6	5.6	7.9	3.3	

*Calculated average length of fish for each age class at a specific age in their life history.



Figure 6.--Population characteristics of largemouth bass collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Table 7.--*Mean back-calculated lengths for northern pike collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
2	37.0	1		31.4	39.8				
3	47.8	1		28.9	34.3	45.7			
4	66.3	5		32.8	42.1	55.1	59.9		
5	67.4	2		33.8	40.5	46.9	60.3	65.0	
6	74.6	2		34.6	43.0	53.1	60.0	63.5	66.2
Mean length of fish at each age				32.3	39.9	50.2	60.1	64.2	66.2
Annual mean growth increment				7.6	10.3	9.9	4.1	2.0	

*Calculated average length of fish for each age class at a specific age in their life history.



Figure 7.--Population characteristics of northern pike collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Table 8.--*Mean back-calculated lengths for yellow perch collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
3	17.0	2		15.6	16.2	17.0			
4	19.6	6		16.0	17.0	18.0	18.7		
5	19.9	4		16.0	17.0	17.9	18.8	19.3	
Mean length of fish at each age				15.9	16.8	17.6	18.8	19.3	
Annual mean growth increment				.9	.8	1.2	.5		

*Calculated average length of fish for each age class at a specific age in their life history.



Figure 8.--Population characteristics of yellow perch collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Table 9.--*Mean back-calculated lengths for rock bass collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
4	12.5	3	11.4	12.0	12.8	13.6			
5	17.0	3	11.1	11.8	12.6	13.3	14.1		
6	15.1	3	11.1	11.7	12.5	13.1	13.7	14.3	
Mean length of fish at each age			11.2	11.8	12.6	13.3	13.9	14.3	
Annual mean growth increment			0.6	0.8	0.7	0.6	0.4		

*Calculated average length of fish for each age class at a specific age in their life history.



Figure 9.--Population characteristics of rock bass collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Table 10.--*Mean back-calculated lengths for black crappie collected in August 1982*

[cm, centimeters]

Age class (years)	Mean length of fish for age class (cm)	Number of specimens	Back-calculated lengths* for indicated age (cm)						
			age (years)						
			1	2	3	4	5	6	7
2	11.4	1	15.2	16.7					
5	23.1	2	15.5	16.9					
6	23.3	2	16.3	17.7	19.2	20.9	21.8		22.6
Mean length of fish at each age			15.7	17.1	18.7	19.7	20.6		22.6
Annual mean growth increment			1.4	1.6	1.0	.9	2.0		

*Calculated average length of fish for each age class at a specific age in their life history.


A
Number of fish = 5
PERCENT
LENGTH AT MIDDLE OF INTERVAL, IN CENTIMETERS
B
Number of fish = 5
PERCENT
AGE, IN YEARS
C
Number of fish = 5
Regression equation Log weight = -2.87 + 3.51 × Log length
R² = .99
DATA POINTS--Each symbol represents the corresponding number of data points
o = 1
Δ = 2
LOG WEIGHT, IN GRAMS
LOG LENGTH, IN CENTIMETERS

Figure 10.--Population characteristics of black crappie collected in August 1982. *A*. Relative abundance by length. *B*. Relative abundance by age. *C*. Relation of length to weight.

Table 11.--*Stomach analyses of fish collected, August 31-September 2, 1982*

Sample size	Species	Number of fish	Terrestrial insects	Odonata	Gastropoda	Diptera larvae	Chironomid larvae	Amphipoda	Daphnia
4	largemouth bass--	2	0	2	0	1	0	14	55
4	bluegill---------	0	0	0	0	2	22	66	0
4	rock bass--------	1	2	5	0	0	1	0	0
1	black crappie---	1	1	0	1	0	0	0	0
2	yellow perch----	0	0	1	0	0	0	0	0

REFERENCES CITED

Bagenal, Timothy, ed., 1978, Methods for assessment of fish production in fresh waters, International Biological Programme Handbook Number 3, (3d ed.): Oxford, Blackwell Scientific Publications, 365 p.

Carlander, K.D., 1977, Handbook of freshwater fishery biology: Ames, Iowa State University Press, v. II, 431 p.

LaBaugh, J.W., Groschen, G.E., Winter, T.C., 1981, Limnological and geochemical survey of Williams Lake, Hubbard County, Minnesota: U.S. Geological Survey Water-Resources Investigations 81-41, 38 p.

Ricker, W.E., 1975, Computation and interpretation of the biological statistics of fish populations: Ottawa, Fisheries Research Board of Canada, Bulletin 191, 382 p.

Siegel, D.I., and Winter, T.C., 1980, Hydrologic setting of Williams Lake, Hubbard County, Minnesota: U.S. Geological Survey Open-File Report 80-403, 56 p.

: U.S. GOVERNMENT PRINTING OFFICE: 1985—677-116/45062 REGION NO. 8